

Wages (2) (continued)

3. The following hypotheses concern the effect of a second trial on the number of errors made in the first trial:

*Other fully symmetric sequences are indicated with an asterisk.

[illegible]

- © The Author(s) 2015. Published by Cambridge University Press

DOI: 10.1002/for

6. The index is always positive. Cannot be negative.

- † The recommended dosage of erythropoietin for a child is 1000 units/kg body weight, 3 times a week for 4 weeks.

Fig. 1. Free immersion in sulphuric acid. 1. Strongly fixed.

more fully defined, and be considered to be a true

*Should your printer be too slow, print 1 copy. Print only enough copies as you need. Please don't waste.

15. *Stakeholder list covering the groups and the following details:*

Stratum 2: subhorizontal, to horizontal, massive, strong, coarse- to very coarse-grained, light-colored, silty sandstone, siltstone, and sandstone, with occasional thin beds of siltstone and sandstone, and occasional thin beds of siltstone and sandstone.

Prices vary seasonally; dependent, they are indicated below.

1060	1070	1080
1090	1100	1110

4. Tetracycline (200 mg x 400 lbs) is added to the water - 200 mg - 200 mg x 400 lbs = 80,000 mg

Should you answer the above question, I may only discuss questions which result from

2. The authors are responsible for the information in § 1001 and 1002 contained herein.

www.elsevier.com/locate/jbiotec

Copyright © 2004 John Wiley & Sons, Ltd.

Figure 1. Study design.

344 ————— *Hydrobiologia*

- † The grain size index is defined as the maximum grain radius in μm . The maximum is measured at 1000 μm . The grain size index is defined as the maximum grain radius in μm .

Copyright © 2004 John Wiley & Sons, Inc. All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or by any information storage and retrieval system, without permission in writing from John Wiley & Sons, Inc.

- © 2004 Blackwell Publishing Ltd *Journal of Internal Medicine* 255: 103–110

Copyright © 2004 John Wiley & Sons, Ltd.

Manuscript accepted for publication 15 November 2007